

# Fitting $K_V$ potassium channels into the $PIP_2$ puzzle: Hille group connects dots between illustrious HH groups

Donald W. Hilgemann

Department of Physiology, UT Southwestern Medical Center at Dallas, Dallas, TX 75235

In the August 2012 issue of the *JGP*, the Bertil Hille group continues its exploration of the physiological significance and cell signaling roles of  $PIP_2$  interactions with potassium channels. As we have come to expect, the group brings surprises and deep insights. In their article (Kruse et al., 2012), they have tackled the question of which voltage-gated potassium ( $K_V$ ) channels may be affected by, and therefore regulated by, changes of  $PIP_2$  that can occur in intact cells. The work brings equally clarity and sobriety to an area of ion channel regulation that has seen false starts. Whereas most, if not all, inward rectifier potassium channels are strongly  $PIP_2$  dependent, the situation with  $K_V$  channels is very different.  $PIP_2$  effects on  $K_V$  channels can be large, small, variable, or absent, and  $PIP_2$  can evidently modify primarily activation/deactivation parameters, inactivation, or channel availability per se in different cases.

Two cases can illustrate the nature of the issues at hand for  $K_V$  channels. First, many  $K_V$  channels, like other channels, are functionally labile in excised patches. Knowing that  $PIP_2$  activates many ion channels, we tested many years ago whether the notoriously labile  $K_V2.1$  channels (Zühlke et al., 1994) might be stabilized in excised patches by incorporating  $PIP_2$ . In our hands, the  $K_V2.1$  channels were completely rescued from run-down and remained stable for unlimited recording periods (Hilgemann et al., 2001). Second, as just noted,  $K_V$  channel inactivation can be drastically “ $PIP_2$  dependent” in excised patches. When recording  $K_V1.1$  or  $K_V3.4$  channel currents in excised patches, the application of  $PIP_2$  micelles to the cytoplasmic side of patches can rapidly abolish inactivation (Oliver et al., 2004). One explanation for the inactivation block is that  $PIP_2$  interacts with the positively charged inactivation domain of  $K_V$  channels. Similar to suggestions for Na/Ca exchangers (Hilgemann and Ball, 1996; He et al., 2000), positively charged inactivation domains might be able to reach and bind to  $PIP_2$  in the plasmalemma, thereby preventing them from causing inactivation. Both of these cases, and others, suggested to many of us that  $PIP_2$  would turn out to be a very important regulator of  $K_V$  channels.

Enter now the Hille group, which is focused on cell signaling roles of  $PIP_2$  that can be demonstrated to occur in intact cells. Previous efforts of this group (Suh and Hille, 2002; Suh et al., 2004; Falkenburger et al., 2010) and the David Brown group (Hughes et al., 2007) established by far the best models yet of physiological regulation of ion channels by  $PIP_2$ , namely by classical pathways involving Gq-coupled receptors whose activation can lead to  $PIP_2$  depletion and subsequent M-type ( $K_V7.2/7.3$ ) potassium channel inhibition. These groups established beyond any reasonable doubt that this pathway controls the firing pattern of these neurons that in turn controls the subsequent release of catecholamines. In addition to modulating M-currents,  $PIP_2$  depletion consequent to muscarinic activation also clearly inhibits N-type Ca channel currents (Gamper et al., 2004). There are in fact very few other examples in which  $PIP_2$  has clearly been shown to play a second messenger role in intact primary cells without receptor overexpression. One very important case, still not resolved clearly, is light-induced depolarization of the *Drosophila melanogaster* rhabdomere, which may be related to  $PIP_2$  depletion or generation of DAG during light-induced phospholipase C (PLC) activation (Huang et al., 2010).

To probe whether  $K_V$  potassium channels (and other channels) can be regulated by physiological  $PIP_2$  changes, the Hille group used several powerful molecular biological tools to rapidly deplete  $PIP_2$  and simultaneously monitored the  $PIP_2$  changes that occurred by FRET between two fluorescent  $PIP_2$ -binding proteins. To deplete  $PIP_2$ , the group used overexpressed muscarinic receptors that couple to endogenous PLCs,  $PIP_2$ -selective phosphatases that are voltage dependent in the sense of being activated by depolarization, and they used  $PIP/PIP_2$ -selective phosphatases that can be brought to the cell surface rapidly “on order” by applying rapamycin. The outcomes were clear and surprising. The same  $K_V$  channels that many of us expected to be  $PIP_2$  sensitive were unaffected by  $PIP_2$  depletion. Thus, large  $PIP_2$  changes in the cell surface of intact cells will not modulate the function of these  $K_V$  channels.

Correspondence to Donald W. Hilgemann:  
Donald.hilgemann@utsouthwestern.edu

There are at least three reasons in my opinion for the striking disparities between expectations from previous experiments and the results of the Hille group. First, the application of PIP<sub>2</sub> micelles to excised patches may generate many physiologically irrelevant results: PIP<sub>2</sub> micelles may interact with positively charged protein domains in ways that never occur physiologically; PIP<sub>2</sub> micelles may extract nonintegral PIP<sub>2</sub>-binding proteins from membranes; and PIP<sub>2</sub> may incorporate to a physiologically irrelevant extent in the cytoplasmic membrane monolayer, thereby generating unphysiological protein confirmations.

Second, PIP<sub>2</sub> may modulate K<sub>V</sub> channels in a constitutive fashion (i.e., in a quasi-structural sense), whereby high affinity PIP<sub>2</sub> interactions might never be subject to regulation by plasmalemmal PIP<sub>2</sub> changes that can occur physiologically. In our experience, this appeared to be the case more often than not, leading us to suggest that PIP<sub>2</sub> is usually acting in a constitutive manner to activate channels and transporters when they are resident in the surface membrane of cells (Hilgemann et al., 2001). Lack of PIP<sub>2</sub>, and perhaps other membrane composition differences, would keep these same channels and transporters inactive when trafficking on internal membranes (Hilgemann et al., 2001). Consistent with this explanation, the Logothetis group has recently described that changes of the gating properties of Shaker-type K<sub>V</sub>1.1 channels, which occur on excision of membrane patches, are largely explained by a loss of PIP<sub>2</sub> (Rodriguez-Menchaca et al., 2012). Because this group used multiple experimental approaches, besides the application of PIP<sub>2</sub> micelles, it would be hard to deny a constitutive function of PIP<sub>2</sub> at these K<sub>V</sub> channels. That said, the Logothetis group does find substantial changes of Shaker potassium channel gating when they induce PIP<sub>2</sub> changes in intact cells. Potential reasons for differences between results of the two groups include both the cell types used and the methods to change PIP<sub>2</sub>.

Third, another perspective says that “PIP<sub>2</sub> is not alone.” The surface membrane is more “anionic” than other membranes of cells, owing to the presence of multiple anionic lipid species that may (or may not) be subject to regulation (Yeung et al., 2008). Our experience is that transporters (Collins et al., 1992) and ion channels are usually not highly selective for PIP<sub>2</sub> versus other anionic lipids (Hilgemann et al., 2001). Thus, biochemically labile anionic lipids besides PIP<sub>2</sub> may activate K<sub>V</sub> channels in intact cells, in concert with PIP<sub>2</sub> or as substitutes for PIP<sub>2</sub>. Phosphatidylserine and phosphatidate are obvious candidates. Whereas phosphatidylserine in the cytoplasmic monolayer may change with alterations of membrane asymmetry that can occur in pathological cell circumstances after membrane patch excision (Hilgemann and Collins, 1992), phosphatidate can alter dramatically with cell signaling changes

(e.g., Lopez et al., 2012). Decades have passed since these issues first arose, and yet we still need to know whether metabolism of lipids besides PIP<sub>2</sub> might underlie the rundown of different ion channels upon excision of membrane patches from cells.

There are many more cases where the methods used by the Hille group (Kruse et al., 2012) can tell us whether channels and transporters might be regulated by changes of surface membrane PIP<sub>2</sub> during cell signaling, starting with a long list of TRP channels that appear to be lipid sensitive (Rohacs, 2009). Ca-activated “BK” potassium channels also belong in this list. In our hands, BK potassium channels were completely insensitive to PIP<sub>2</sub> changes, using patches from tracheal smooth muscle cells, thereby verifying to us that PIP<sub>2</sub> sensitivity was specific to specific channel types (Hilgemann and Ball, 1996). However, it turned out that the cell-specific expression of β-1 subunits of these channels confers PIP<sub>2</sub> sensitivity (Vaithianathan et al., 2008). The involvement of subunits in establishing PIP<sub>2</sub> sensitivity is clearly consistent with the idea that PIP<sub>2</sub> signaling to these channels will not only be “constitutive” in nature but also occurs in a cell environment-critical fashion. Cardiac Na/Ca exchangers and K<sub>ATP</sub> channels were the first two ion transport systems that my group found to be PIP<sub>2</sub> sensitive. Both cardiac Na/Ca exchangers and K<sub>ATP</sub> channels were strongly activated by the ATP-dependent generation of PIP<sub>2</sub> in membrane patches, as well as by exogenous PIP<sub>2</sub> (Hilgemann and Ball, 1996). Nevertheless, the physiological roles that PIP<sub>2</sub> plays in both systems remain rather enigmatic (Hilgemann, 2007).

These examples illustrate important gaps in our knowledge about the role of PIP<sub>2</sub> for individual channels and transporters, gaps that can be filled by using the tools used by the Hille group. These tools can also be used to address larger questions about PIP<sub>2</sub> functions in cells. Besides its complex roles in cytoskeleton regulation and classical endocytic processes, PIP<sub>2</sub> appears to drive the formation of membrane domains, membrane budding, and membrane turnover (Fine et al., 2011; Hilgemann and Fine, 2011; Lariccia et al., 2011). In this context, the present studies of K<sub>V</sub> channel regulation by PIP<sub>2</sub> reflect a convergence of the work of two great “H&H” research teams of the early 1950s. The first team was of course Hodgkin and Huxley, whose analysis of voltage-gated ion channels is known to every reader of this journal. The second team, less well known to the readers of this journal, was Hokin and Hokin, who described in the same years that phospholipids become phosphorylated in response to autonomic hormones, specifically that acetylcholine activates phosphorylation of lipids in secretory cells (Hokin and Hokin, 1953). Hokin and Hokin’s results represent the activation of synthesis pathways for multiple phospholipids, including PIP<sub>2</sub>, other phosphoinositides, and

phosphatidate, by the same general pathways that activate PIP<sub>2</sub> breakdown by PLCs (Hokin, 1996). As surprising as it may be, these pathways remain poorly understood because turnover of these lipids is connected to membrane turnover, so that an understanding of the underlying mechanisms becomes highly cell biological. For certain, phosphatidylinositol must be continuously provided to the surface membrane from intracellular organelles, where it is generated (Kim et al., 2011). There are also indications that PI4P is not generated at the cell surface but is transferred from other membrane sources (Yaradanakul et al., 2007; Balla et al., 2008), whereas PIP<sub>2</sub> is lost from the surface membrane during endocytosis (Doherty and McMahon, 2009). These issues must ultimately be understood to understand how PIP<sub>2</sub> metabolism controls ion channels in intact cells. To no surprise, Bertil Hille's group is now investigating these more cell biological issues (Dickson, E.J., G.R.V. Hammond, and B. Hille. 2012. Biophysical Society 56th Annual Meeting. Abstr. 1681) in addition to PIP<sub>2</sub>-channel interactions. With new light-activated enzymatic probes moving onto their stage (Idevall-Hagren et al., 2012), we may soon expect new insights into both the biophysical and cell biological basis of K<sub>V</sub> channel regulation by PIP<sub>2</sub>.

## REFERENCES

- Balla, A., Y.J. Kim, P. Varnai, Z. Szentpetery, Z. Knight, K.M. Shokat, and T. Balla. 2008. Maintenance of hormone-sensitive phosphoinositide pools in the plasma membrane requires phosphatidylinositol 4-kinase IIIalpha. *Mol. Biol. Cell.* 19:711–721. <http://dx.doi.org/10.1091/mbc.E07-07-0713>
- Collins, A., A.V. Somlyo, and D.W. Hilgemann. 1992. The giant cardiac membrane patch method: stimulation of outward Na(+)-Ca2+ exchange current by MgATP. *J. Physiol.* 454:27–57.
- Doherty, G.J., and H.T. McMahon. 2009. Mechanisms of endocytosis. *Annu. Rev. Biochem.* 78:857–902. <http://dx.doi.org/10.1146/annurev.biochem.78.081307.110540>
- Falkenburger, B.H., J.B. Jensen, and B. Hille. 2010. Kinetics of PIP<sub>2</sub> metabolism and KCNQ2/3 channel regulation studied with a voltage-sensitive phosphatase in living cells. *J. Gen. Physiol.* 135:99–114. <http://dx.doi.org/10.1085/jgp.200910345>
- Fine, M., M.C. Llaguno, V. Lariccia, M.J. Lin, A. Yaradanakul, and D.W. Hilgemann. 2011. Massive endocytosis driven by lipid forces originating in the outer plasmalemmal monolayer: a new approach to membrane recycling and lipid domains. *J. Gen. Physiol.* 137:137–154. <http://dx.doi.org/10.1085/jgp.201010469>
- Gamper, N., V. Reznikov, Y. Yamada, J. Yang, and M.S. Shapiro. 2004. Phosphatidylinositol 4,5-bisphosphate signals underlie receptor-specific Gq/11-mediated modulation of N-type Ca2+ channels. *J. Neurosci.* 24:10980–10992. <http://dx.doi.org/10.1523/JNEUROSCI.3869-04.2004>
- He, Z., S. Feng, Q. Tong, D.W. Hilgemann, and K.D. Philipson. 2000. Interaction of PIP(2) with the XIP region of the cardiac Na/Ca exchanger. *Am. J. Physiol. Cell Physiol.* 278:C661–C666.
- Hilgemann, D.W. 2007. On the physiological roles of PIP(2) at cardiac Na+ Ca2+ exchangers and K(ATP) channels: a long journey from membrane biophysics into cell biology. *J. Physiol.* 582:903–909. <http://dx.doi.org/10.1113/jphysiol.2007.132746>
- Hilgemann, D.W., and R. Ball. 1996. Regulation of cardiac Na+,Ca2+ exchange and KATP potassium channels by PIP2. *Science.* 273:956–959. <http://dx.doi.org/10.1126/science.273.5277.956>
- Hilgemann, D.W., and A. Collins. 1992. Mechanism of cardiac Na(+)-Ca2+ exchange current stimulation by MgATP: possible involvement of aminophospholipid translocase. *J. Physiol.* 454:59–82.
- Hilgemann, D.W., and M. Fine. 2011. Mechanistic analysis of massive endocytosis in relation to functionally defined surface membrane domains. *J. Gen. Physiol.* 137:155–172. <http://dx.doi.org/10.1085/jgp.201010470>
- Hilgemann, D.W., S. Feng, and C. Nasuhoglu. 2001. The complex and intriguing lives of PIP2 with ion channels and transporters. *Sci. STKE.* 2001:re19. <http://dx.doi.org/10.1126/stke.2001.111.re19>
- Hokin, L.E. 1996. History of phosphoinositide research. *Subcell. Biochem.* 26:1–41. [http://dx.doi.org/10.1007/978-1-4613-0343-5\\_1](http://dx.doi.org/10.1007/978-1-4613-0343-5_1)
- Hokin, M.R., and L.E. Hokin. 1953. Enzyme secretion and the incorporation of P32 into phospholipides of pancreas slices. *J. Biol. Chem.* 203:967–977.
- Huang, J., C.H. Liu, S.A. Hughes, M. Postma, C.J. Schwiening, and R.C. Hardie. 2010. Activation of TRP channels by protons and phosphoinositide depletion in Drosophila photoreceptors. *Curr. Biol.* 20:189–197. <http://dx.doi.org/10.1016/j.cub.2009.12.019>
- Hughes, S., S.J. Marsh, A. Tinker, and D.A. Brown. 2007. PIP(2)-dependent inhibition of M-type (Kv7.2/7.3) potassium channels: direct on-line assessment of PIP(2) depletion by Gq-coupled receptors in single living neurons. *Pflugers Arch.* 455:115–124. <http://dx.doi.org/10.1007/s00424-007-0259-6>
- Idevall-Hagren, O., E.J. Dickson, B. Hille, D.K. Toomre, and P. De Camilli. 2012. Optogenetic control of phosphoinositide metabolism. *Proc. Natl. Acad. Sci. USA.* In press.
- Kim, Y.J., M.L. Guzman-Hernandez, and T. Balla. 2011. A highly dynamic ER-derived phosphatidylinositol-synthesizing organelle supplies phosphoinositides to cellular membranes. *Dev. Cell.* 21:813–824. <http://dx.doi.org/10.1016/j.devcel.2011.09.005>
- Kruse, M., G.R.V. Hammond, and B. Hille. 2012. Regulation of voltage-gated potassium channels by PI(4,5)P<sub>2</sub>. *J. Gen. Physiol.* 140:189–205. <http://dx.doi.org/10.1085/jgp.201210806>
- Lariccia, V., M. Fine, S. Magi, M.J. Lin, A. Yaradanakul, M.C. Llaguno, and D.W. Hilgemann. 2011. Massive calcium-activated endocytosis without involvement of classical endocytic proteins. *J. Gen. Physiol.* 137:111–132. <http://dx.doi.org/10.1085/jgp.201010468>
- Lopez, C.I., L.E. Pelletán, L. Suhaiman, G.A. De Blas, N. Vitale, L.S. Mayorga, and S.A. Belmonte. 2012. Diacylglycerol stimulates acrosomal exocytosis by feeding into a PKC- and PLD1-dependent positive loop that continuously supplies phosphatidylinositol 4,5-bisphosphate. *Biochim. Biophys. Acta.* 1821:1186–1199. <http://dx.doi.org/10.1016/j.bbali.2012.05.001>
- Oliver, D., C.C. Lien, M. Soom, T. Baukowitz, P. Jonas, and B. Fakler. 2004. Functional conversion between A-type and delayed rectifier K+ channels by membrane lipids. *Science.* 304:265–270. <http://dx.doi.org/10.1126/science.1094113>
- Rodríguez-Menchaca, A.A., S.K. Adney, Q.-Y. Tang, X.-Y. Meng, A. Rosenhouse-Dantsker, M. Cui, and D.E. Logothetis. 2012. PIP<sub>2</sub> controls voltage-sensor movement and pore opening of Kv channels through the S4-S5 linker. *Proc. Natl. Acad. Sci. USA.* In press. <http://dx.doi.org/10.1073/pnas.1207901109>
- Rohacs, T. 2009. Phosphoinositide regulation of non-canonical transient receptor potential channels. *Cell Calcium.* 45:554–565. <http://dx.doi.org/10.1016/j.ceca.2009.03.011>
- Suh, B.C., and B. Hille. 2002. Recovery from muscarinic modulation of M current channels requires phosphatidylinositol 4,5-bisphosphate synthesis. *Neuron.* 35:507–520. [http://dx.doi.org/10.1016/S0896-6273\(02\)00790-0](http://dx.doi.org/10.1016/S0896-6273(02)00790-0)

- Suh, B.C., L.F. Horowitz, W. Hirdes, K. Mackie, and B. Hille. 2004. Regulation of KCNQ2/KCNQ3 current by G protein cycling: the kinetics of receptor-mediated signaling by  $G_q$ . *J. Gen. Physiol.* 123:663–683. <http://dx.doi.org/10.1085/jgp.200409029>
- Vaithianathan, T., A. Bukiya, J. Liu, P. Liu, M. Asuncion-Chin, Z. Fan, and A. Dopico. 2008. Direct regulation of BK channels by phosphatidylinositol 4,5-bisphosphate as a novel signaling pathway. *J. Gen. Physiol.* 132:13–28. <http://dx.doi.org/10.1085/jgp.200709913>
- Yaradanakul, A., S. Feng, C. Shen, V. Lariccia, M.-J. Lin, J. Yang, T.M. Kang, P. Dong, H.L. Yin, J.P. Albanesi, and D.W. Hilgemann. 2007. Dual control of cardiac  $Na^+$   $Ca^{2+}$  exchange by PIP(2): electrophysiological analysis of direct and indirect mechanisms. *J. Physiol.* 582:991–1010. <http://dx.doi.org/10.1113/jphysiol.2007.132712>
- Yeung, T., G.E. Gilbert, J. Shi, J. Silvius, A. Kapus, and S. Grinstein. 2008. Membrane phosphatidylserine regulates surface charge and protein localization. *Science*. 319:210–213. <http://dx.doi.org/10.1126/science.1152066>
- Zühlke, R.D., H.J. Zhang, and R.H. Joho. 1994. Role of an invariant cysteine in gating and ion permeation of the voltage-sensitive  $K^+$  channel Kv2.1. *Receptors Channels*. 2:237–248.